

Published by the British
Herpetological Society**Appendix 1. Literature containing environmental parameters used in the analysis presented in Table 3.**

- Afonso, L.G. & Eterovick, P.C. (2007). Microhabitat choice and differential use by anurans in forest streams in southeastern Brazil. *Journal of Natural History* 41, 937–948.
- Banks, B. & Beebee, T.J. (1987). Factors influencing breeding site choice by the pioneering amphibian *Bufo calamita*. *Ecography* 10, 14–21.
- Baumgartner, N., Waringer, A. & Waringer, J. (1999). Hydraulic microdistribution patterns of larval fire salamanders (*Salamandra salamandra salamandra*) in the Weidlingbach near Vienna, Austria. *Freshwater Biology* 41, 31–41.
- Bei, Y., Meng, S., Li, G., Xie, W., et al. (2012). First record of nest site and egg guarding in the caecilian *Ichthyophis bannanicus* (Amphibia: Gymnophiona: Ichthyophiidae). *Journal of Natural History* 46, 859–865.
- Bernardes, M., Rödder, D., Nguyen, T.T., Pham, C.T., et al. (2013). Habitat characterization and potential distribution of *Tylototriton vietnamensis* in northern Vietnam. *Journal of Natural History* 47, 1161–1175.
- Blackburn, D.C. & Moreau, C.S. (2006). Ontogenetic diet change in the arthroleptid frog *Schoutedenella xenodactyloides*. *Journal of Herpetology* 40, 388–394.
- Boistel, R., Grosjean, S. & Lötters, S. (2005). Tadpole of *Atelopus franciscus* from French Guyana, with comments on other larvae of the genus (Anura: Bufonidae). *Journal of Herpetology* 39, 148–153.
- Born, M., Bongers, F., Poelman, E.H. & Sterck, F.J. (2010). Dry-season retreat and dietary shift of the dart-poison frog *Dendrobates tinctorius* (Anura: Dendrobatidae). *Phylomedusa* 9, 37–52.
- Caldwell, J.P. & Araújo, M.C. (1998). Cannibalistic Interactions Resulting from Indiscriminate Predatory Behavior in Tadpoles of Poison Frogs (Anura: Dendrobatidae). *Biotropica* 30, 92–103.
- Coloma, L.A. & Lötters, S. (1996). The tadpole of *Atelopus balios* (Anura: Bufonidae) from the Pacific lowlands of Ecuador. *Herpetologica* 52, 66–70.
- Dasgupta, R. (1996). Feeding ecology of the adult Himalayan Salamander *Tylototriton verrucosus* Anderson, 1871. *Herpetozoa* 9, 19–29.
- Dole, J.W. (1965). Summer movements of adult leopard frogs, *Rana pipiens* Schreber, in northern Michigan. *Ecology* 46, 236–255.
- Dole, J.W. & Durant, P. (1974). Movements and seasonal activity of *Atelopus oxyrhynchus* (Anura: Atelopodidae) in a Venezuelan cloud forest. *Copeia* 1974, 230–235.
- Freda, J. & Dunson, W.A. (1985). Field and laboratory studies of ion balance and growth rates of ranid tadpoles chronically exposed to low pH. *Copeia* 1985, 415–423.
- Freda, J. & Dunson, W.A. (1986). Effects of low pH and other chemical variables on the local distribution of amphibians. *Copeia* 1986, 454–466.
- Gaborieau, O. & Measey, G.J. (2004). Termitivore or detritivore? A quantitative investigation into the diet of the East African caecilian *Boulengerula taitanus* (Amphibia: Gymnophiona: Caeciliidae). *Animal Biology* 54, 45–56.
- Gong, D. & Mu, M. (2008). Behavioral Observations and Descriptions of the Endangered Knobby Newt *Tylototriton wenxianensis* and Their Application in Conservation. *Asiatic Herpetological Research* 11, 31–38.
- Grant, R.A., Chadwick, E.A. & Halliday, T. (2009). The lunar cycle: a cue for amphibian reproductive phenology?. *Animal Behaviour* 78, 349–357.
- Horne, M.T. & Dunson, W.A. (1994). Behavioral and physiological responses of the terrestrial life stages of the Jefferson salamander, *Ambystoma jeffersonianum*, to low soil pH. *Archives of environmental contamination and toxicology* 27, 232–238.
- Kaefer, I.L., Both, C. & Cechin, S.Z. (2009). Breeding biology of the rapids frog *Limnonectes macroglossa* (Anura: Cycloramphidae) in southern Brazil. *Journal of Natural History* 43, 1195–1206.
- Karraker, N.E., Richards, C.L. & Ross, H.L. (2006). Reproductive ecology of *Atelopus zeteki* and comparisons to other members of the genus. *Herpetological Review* 37, 284–287.
- Kluber, M.R., Olson, D.H. & Puettmann, K.J. (2009). Downed wood microclimates and their potential impact on plethodontid salamander habitat in the Oregon Coast Range. *Northwest Science* 83, 25–34.
- Kopp, K., & Eterovick, P.C. (2006). Factors influencing spatial and temporal structure of frog assemblages at ponds in southeastern Brazil. *Journal of Natural History*, 40, 1813–1830.
- Kovacs, E.H., Sas, I., Covaci-Marcov, S.D., Hartel, T., et al. (2007). Seasonal variation in the diet of a population of *Hyla arborea* from Romania. *Amphibia-Reptilia* 28, 28–34.
- Kupfer, A., Nabhitabhata, J. & Himstedt, W. (2005). Life history of amphibians in the seasonal tropics: habitat, community and population ecology of a caecilian (genus *Ichthyophis*). *Journal of Zoology* 266, 237–247.
- Lescure, J. (1981). Contribution à l'étude des Amphibiens de Guyane française IX. Le têtard gastromyzophage d'*Atelopus flavescens* Dumeril et Bibron (Anura, Bufonidae). *Amphibia-Reptilia* 2, 209–215.
- Leuven, R.S.E.W., Den Hartog, C., Christiaans, M.M.C. & Heijligers, W.H.C. (1986). Effects of water acidification on the distribution pattern and the reproductive success of amphibians. *Experientia* 42, 495–503.
- Lillywhite, H.B., Licht, P. & Chelgren, P. (1973). The role of behavioral thermoregulation in the growth energetics of

- the toad, *Bufo boreas*. *Ecology* 54, 375–383.
- Lindquist, E.D. & Hetherington, T.E. (1998). Tadpoles and juveniles of the Panamanian golden frog, *Atelopus zeteki* (Bufonidae), with information on development of coloration and patterning. *Herpetologica* 54, 370–376.
- Lötters, S., Schick, S., Scheelke, K., Teege, P., et al. (2004). Bio-sketches and partitioning of sympatric reed frogs, genus *Hyperolius* (Amphibia; Hyperoliidae), in two humid tropical African forest regions. *Journal of Natural History* 38, 1969–1997.
- Malonza, P.K., & Measey, G.J. (2005). Life history of an African caecilian: *Boulengerula taitanus* Loveridge 1935 (Amphibia Gymnophiona Caeciliidae). *Tropical Zoology* 18, 49–66.
- Measey, G.J. (2004). Are caecilians rare? An East African perspective. *Journal of East African Natural History* 93, 1–21.
- Measey, G.J. & Barot, S. (2006). Evidence of seasonal migration in a tropical subterranean vertebrate. *Journal of Zoology* 269, 29–37.
- Measey, G.J. & Di-Bernardo, M. (2003). Estimating juvenile abundance in a population of the semiaquatic caecilian, *Chttonerpeton indistinctum* (Amphibia: Gymnophiona: Typhlonectidae), in southern Brazil. *Journal of Herpetology* 37, 371–373.
- Measey, G.J., Hinkel, H.H., Dumbo, B. & Fischer, E. (2011). Rediscovery of *Boulengerula fischeri*, with notes on its morphology and habitat. *African Journal of Herpetology* 60, 47–59.
- Mercurio, V. & Andreone, F. (2006). The tadpoles of *Scaphiophryne gottlebei* (Microhylidae: Scaphiophryinae) and *Mantella expectata* (Mantellidae: Mantellinae) from Isalo Massif, south-central Madagascar. *Alytes* 23, 81–95.
- Miller, N. (1909). The American Toad (*Bufo Lentiginosus americanus*, LeConte) A Study in Dynamic Biology. *The American Naturalist* 43, 641–668.
- Navas, C.A. (1996). Implications of microhabitat selection and patterns of activity on the thermal ecology of high elevation neotropical anurans. *Oecologia*, 108, 617–626.
- Navas, C.A. (1996). Thermal dependency of field locomotor and vocal performance of high-elevation anurans in the tropical Andes. *Journal of Herpetology* 30, 478–487.
- Nishikawa, K., Matsui, M. & Nguyen, T.T. (2013). A New Species of *Tylototriton* from Northern Vietnam (Amphibia: Urodela: Salamandridae). *Current Herpetology* 32, 34–49.
- Noland, R. & Ultsch, G.R. (1981). The roles of temperature and dissolved oxygen in microhabitat selection by the tadpoles of a frog (*Rana pipiens*) and a toad (*Bufo terrestris*). *Copeia* 1981, 645–652.
- Oommen, O.V., Measey, G.J., Gower, D.J. & Wilkinson, M. (2000). Distribution and abundance of the caecilian *Gegeneophis ramaswamii* (Amphibia: Gymnophiona) in southern Kerala. *Current Science-Bangalore* 79, 1386–1389.
- Phimmachak, S., Stuart, B.L. & Sivongxay, N. (2012). Distribution, natural history, and conservation of the Lao newt (*Laotriton laoensis*) (Caudata: Salamandridae). *Journal of Herpetology* 46, 120–128.
- Picker, M.D. & de Villiers, A.L. (1989). The distribution and conservation status of *Xenopus gilli* (Anura: Pipidae). *Biological Conservation* 49, 169–183.
- Pough, F.H. (1974). Natural daily temperature acclimation of eastern red efts *Notophthalmus v. viridescens* (Rafinesque) (Amphibia: Caudata). *Comparative Biochemistry and Physiology Part A: Physiology* 47, 71–78.
- Pough, F.H. & Wilson, R.E. (1977). Acid precipitation and reproductive success of *Ambystoma* salamanders. *Water, Air, and Soil Pollution* 7, 307–316.
- Randrianavelona, R., Rakotonoloy, H., Ratsimbazafy, J. & Jenkins, R.K. (2010). Conservation assessment of the critically endangered frog *Mantella aurantiaca* in Madagascar. *African Journal of Herpetology* 59, 65–78.
- Rittenhouse, T.A. & Semlitsch, R.D. (2007). Postbreeding habitat use of wood frogs in a Missouri oak-hickory forest. *Journal of Herpetology* 41, 645–653.
- Rodrigues, D.J., Uetanabaro, M. & Lopes, F.S. (2007). Breeding biology of *Phyllomedusa azurea* Cope, 1862 and *P. sauvagii* Boulenger, 1882 (Anura) from the Cerrado, Central Brazil. *Journal of Natural History* 41, 1841–1851.
- Seebacher, F. & Alford, R.A. (1999). Movement and microhabitat use of a terrestrial amphibian (*Bufo marinus*) on a tropical island: seasonal variation and environmental correlates. *Journal of Herpetology* 33, 208–214.
- Seymour, R.S., Roberts, J.D., Mitchell, N.J. & Blaylock, A.J. (2000). Influence of environmental oxygen on development and hatching of aquatic eggs of the Australian frog, *Crinia georgiana*. *Physiological and Biochemical Zoology* 73, 501–507.
- Strijbosch, H. (1979). Habitat selection of amphibians during their aquatic phase. *Oikos* 33, 363–372.
- Stuart, B.L., Phimmachak, S., Sivongxay, N. & Robichaud, W.G. (2010). A new species in the *Tylototriton asperrimus* group (Caudata: Salamandridae) from central Laos. *Zootaxa* 2650, 19–32.
- Tapley, B. & Acosta-Galvis, A.R. (2010). Distribution of *Typhlonectes natans* in Colombia, environmental parameters and implications for captive husbandry. *Herpetological Bulletin* 113, 23–29.
- Wachlevski, M., De Souza, P.H., Kopp, K. & Eterovick, P.C. (2008). Microhabitat use and feeding habits of *Crossodactylus bokermanni* Caramaschi & Sazima, 1985 (Anura, Hydromidae) at a site in south-eastern Brazil. *Journal of Natural History* 42, 1421–1434.
- Wyman, R.L. (1988). Soil acidity and moisture and the distribution of amphibians in five forests of southcentral New York. *Copeia* 1988, 394–399.
- Xu, F. & Li, Y. (2013). Oviposition site selection by rice frogs on Taohua Island and the nearby mainland. *The Herpetological Journal* 23, 51–53.