



# Host-pathogen relationships between the chytrid fungus *Batrachochytrium dendrobatidis* and tadpoles of five South American anuran species

María Luz Arellano<sup>1</sup>, Guillermo S. Natale<sup>2</sup>, Pablo G. Grilli<sup>3</sup>, Diego A. Barrasso<sup>4</sup>,  
Mónica M. Steciow<sup>5</sup> & Esteban O. Lavilla<sup>6</sup>

<sup>1</sup>Instituto de Botánica Spegazzini, Facultad de Ciencias Naturales y Museo (FCNyM), Universidad Nacional de La Plata (UNLP), Calle 53 No 477, 1900 La Plata, Buenos Aires, Argentina

<sup>2</sup>CIMA, Departamento de Química, Facultad de Ciencias Exactas (FCE), UNLP, Calle 115 esquina 47, 1900 La Plata, Buenos Aires, Argentina

<sup>3</sup>Cátedra de Ecología General y Recursos Naturales, Universidad Nacional Arturo Jauretche (UNAJ), Av. Calchaquí 6200, 1888 Florencio Varela, Buenos Aires, Argentina

<sup>4</sup>Instituto de Diversidad y Evolución Austral - CONICET. Blvd. Brown 2915, U9120ACD Puerto Madryn, Chubut, Argentina

<sup>5</sup>Instituto de Botánica Spegazzini, FCNyM, UNLP, Calle 53 No 477, 1900 La Plata, Buenos Aires, Argentina

<sup>6</sup>Unidad Ejecutora Lillo, Conicet-Fundación Miguel Lillo, Miguel Lillo 251, 4000 San Miguel de Tucumán, Argentina

The chytrid fungus *Batrachochytrium dendrobatidis* (*Bd*) is one of the most important contributors for the decline of amphibian populations worldwide. Evidence indicates that the harmfulness of *Bd* infection depends on the species and life stage, the fungus strain, the season and environmental factors. In the present paper, we experimentally investigated (i) the susceptibility and sensitivity of five South American tadpole species (*Rhinella fernandezae*, *Scinax squalirostris*, *Hypsiboas pulchellus*, *Leptodactylus latrans* and *Physalaemus fernandezae*) to a foreign *Bd* strain (JEL423), (ii) the response of two populations of *P. fernandezae* to a native *Bd* strain (MLA1), and (iii) the virulence of native and foreign *Bd* isolates on tadpoles of the same species. We also evaluated the relationship between *Bd* infection and the loss of keratinised mouthparts in *P. fernandezae*. We found that all species except *L. latrans* were susceptible to *Bd* infection with lethal consequences, with *R. fernandezae* being the most sensitive species. In *P. fernandezae*, sensitivity to infection depended on population as well as *Bd* strain, although no relationship was found between fungal infection and the loss of keratinised mouthparts. This is the first experimental study on mortality rates of South American tadpoles exposed to *Bd*.

**Key words:** *Batrachochytrium dendrobatidis*, fungal infection, South America, tadpoles

## INTRODUCTION

The pathogenic fungus *Batrachochytrium dendrobatidis* (*Bd*), the etiological agent of chytridiomycosis (Longcore et al., 1999), is recognised as a proximate driver of many severe declines of amphibian populations worldwide (Lips et al., 2006). *Bd* infects keratinising tissue such as mouthparts of larvae and the skin of adults (Berger et al., 1998; Altig, 2007). The complexity of host-pathogen interaction in the *Bd*-amphibian system has been studied extensively among different amphibian species, *Bd* strains, populations and environmental conditions (Searle et al., 2011; Gervasi et al., 2013; Ortiz-Santaliestra et al., 2013; Langhammer et al., 2014; Spitzen-Van Der Sluijs et al., 2014). Whereas some species carry constant infections in nature with little or no evidence of disease outbreaks (Kielgast et al., 2009; Reeder et al., 2012), others suffer significant declines (Ryan et al., 2008; Vredenburg et al., 2010). Sensitivity can also vary among amphibian life stages, and tadpoles of most species often show low sensitivity

to *Bd* infection until metamorphosis (Rachowicz & Briggs, 2007; Symonds et al., 2007; Narayan et al., 2014), probably because infections only occur on the keratinised mouthparts (Berger et al., 1998). As a consequence, they can act as *Bd* reservoirs for the pathogen to persist (Blaustein et al., 2005; Mitchell et al., 2008), despite reports of reduced survival due to infection in a range of species (Blaustein et al., 2005; Garner et al., 2009; Gahl et al., 2012; Paetow et al., 2013; Hanlon & Parris, 2014).

Most studies on *Bd* and its relationship with the anuran host are from the northern hemisphere and Australia (see Voyles et al., 2011). We believe it is essential to generate knowledge about infection in South American amphibians, in order to carry out future conservation programs and to identify key species to prioritise. Therefore, we used a comparative experimental approach to examine host responses to *Bd* infection (susceptibility, sensitivity and loss of keratinised mouthparts) in tadpoles of five South American anuran species: the burrowing toad (*Rhinella fernandezae*), the white-banded treefrog (*Hypsiboas pulchellus*),

Correspondence: María Luz Arellano (mluzarellano@gmail.com)

the striped snouted treefrog (*Scinax squalirostris*), the creole frog (*Leptodactylus latrans*), and the whistling dwarf frog (*Physalaemus fernandezae*). Moreover, we investigated the susceptibility of *P. fernandezae* to *Bd* depending on strain and populations. To our knowledge, this study represents the first experimental bioassays on *Bd* infection of native South American tadpoles.

## METHODS

Tadpoles were collected from natural breeding sites in temporary ponds located at three sites in Buenos Aires province, Argentina. *R. fernandezae* (stages 29–34 [Gosner, 1960]), *H. pulchellus* (30–34), *S. squalirostris* (34–37), and *L. latrans* (33–36) larvae were collected in the outskirts of La Plata [35° S, 57° W] and used for Bioassay 1. *P. fernandezae* larvae (stages 33–37) were collected from La Balandra [34° S, 57° W] and Pinamar (25–35) [37° S, 56° W], and used for Bioassays 2 and 3. Data from individual *P. fernandezae* from La Balandra were also used in Bioassay 1. Upon arrival at the laboratory, all individuals were held at 37°C for 16 h to eliminate any *Bd* (Woodhams et al., 2003); tadpoles were subsequently acclimated for five days at 17°C, and a photoperiod of 16:8 hours of light:dark. After acclimation, tadpoles were placed individually into 500 ml cylindric polypropylene containers with perforated plastic lids and 56 ml of dechlorinated water.

Zoospores we collected from two different *Bd* strains: JEL423 isolated from an adult *Agalychnis lemur* from Panama, and MLA1 isolated from larvae of *Hypsiboas cordobae* from Argentina. Zoospore collection was done by washing three-day-old 1% tryptone agar plates (grown at 23°C) for 1 hour with 4 ml of distilled water over three consecutive days (the same procedure was performed with *Bd*-free agar plates for control groups), obtaining a final suspension of  $4 \times 10^6$  zoospore ml<sup>-1</sup>. A Neubauer chamber was used for zoospore counts.

For *Bd* exposure treatments, we inoculated containers containing 56 ml of dechlorinated water with 4 ml of daily harvested zoospore suspension (*Bd*-free suspension for the control group) for three consecutive days (exposure time), obtaining a final concentration of  $6 \times 10^4$  zoospore ml<sup>-1</sup> in each container. After this period, the water in experimental containers was replaced with fungus-free water every day. Tadpoles were fed liquefied lettuce *ad libitum* and checked daily for mortality counts. Bioassays were ended when mortality was recorded in all exposed individuals, or in 10% of the individuals of the control group. A solution of the anesthetic MS222 (tricaine methane sulfonate) was used to humanely euthanise tadpoles, which were then fixed in 10% formalin. To determine the presence of abnormalities in keratinised mouthparts, we extracted the oral disc for inspection with a stereomicroscope (Wild M3 Heerbrugg). *Bd* presence was identified through direct and histological examination of oral structures following Berger et al. (1999; hematoxylin and eosin staining using a compound optical microscope Hund Wetzlar H600).

Results were assessed considering three criteria: (1) susceptibility, defined as the ability to become infected

with *Bd* (a species was considered susceptible to *Bd* when at least one individual was infected with the pathogen); (2) sensitivity, defined as survival time after *Bd* exposure; (3) mouthpart deformity, as partial or total absence of keratinised mouthparts on the oral disc. To investigate whether anurans species are susceptible to infection by *Bd* and to assess their sensitivity, Bioassay 1 consisted in exposing tadpoles of *R. fernandezae*, *H. pulchellus*, *S. squalirostris*, *L. latrans* and *P. fernandezae* to *Bd* strain JEL423. Results from Bioassay 3 (using *Bd* strain JEL423) performed on *P. fernandezae* were also included in the data analysis. With Bioassay 2, we tested whether different populations of a single species were differentially affected by *Bd*. We used the same experimental design as in Bioassay 1 and compared two *P. fernandezae* populations (La Balandra and Pinamar) to a locally isolated *Bd* strain (MLA1). In Bioassay 3 we exposed tadpoles of *P. fernandezae* from the La Balandra population to MLA1 or to JEL423. Fungal exposures for each trial were conducted simultaneously, and we used 10 exposed and 10 control larvae for each species, population and *Bd* strain.


We used Kaplan-Meier analyses (XLSTAT software version 2013.5.04; Addinsoft) to generate survival curves for species (Bioassay 1), populations (Bioassay 2), and groups exposed to *Bd* strains (Bioassay 3), comparing them using a Log-Rank Test. A contingency table analysis was performed on infection data (*Bd*/no *Bd*) with characteristics of mouthparts (normal/deformed), to determine whether the presence of *Bd* was related to oral disc deformation in *P. fernandezae* (individuals of Bioassays 2 and 3). We considered an oral disc to be deformed when keratinised mouthparts (labial teeth and jaw sheath) were absent (Altig, 2007).

## RESULTS

### Bioassay 1

Nine out of ten *R. fernandezae* individuals, and five out of ten *S. squalirostris* individuals treated with *Bd* died within 24 h of inoculation, and the remainder died on day 2. The other species survived longer, with *P. fernandezae* having the longest survival time (Fig. 1A). No animals in the control groups died except for one tadpole of *L. latrans* on day 5. Treatment and control groups differed significantly for all species, with the following Mean Survival Times (MST) in the treatment groups: *R. fernandezae* ( $p < 0.001$ ) 1.1 d, *S. squalirostris* ( $p < 0.001$ ) 1.5 d, *L. latrans* ( $p = 0.004$ ) 3.4 d, *H. pulchellus* ( $p < 0.001$ ) 3.9 d, and *P. fernandezae* ( $p < 0.001$ ) 6.5 d. We also observed significant differences in survival among all species ( $p < 0.001$ ).

We found no evident oral disc deformations except in *P. fernandezae* (nine out of ten individuals, see Bioassay 2). Four out of the five species tested positive for the presence of *Bd* (direct examination of fresh oral disc surface at 400 × without stain; Fig. 2): *H. pulchellus* (4/10 inoculated individuals), *R. fernandezae* (5/10), *S. squalirostris* (4/10), and *P. fernandezae* (2/10); *L. latrans* tested negative (0/10). The histological analysis of sectioned and stained mouthparts of larvae of all species revealed no evidence of *Bd*.



**Fig. 1.** Relative survival through time. A). Tadpoles of five anuran species exposed to *Batrachochytrium dendrobatidis* (Bd). Hp: *Hypsiboas pulchellus*, Ll: *Leptodactylus latrans*, Pf: *Physalaemus fernandae*, Rf: *Rhinella fernandae*, Sq: *Scinax squalirostris*. B) *P. fernandae* tadpoles from La Balandra and Pinamar exposed to Bd strain MLA1. C) *P. fernandae* tadpoles from La Balandra exposed to Bd strains MLA1 and JEL423. Symbols in lines of control groups show the end of the bioassay for each species. A) Square: Rf and Sq; triangle: Ll; star: Hp; circle: Pf. (no survival in exposed or control individuals but in 10% of Ll control individuals). B) Circle: La Balandra; triangle: Pinamar (no survival in exposed or control individuals). C) Circle: JEL423; triangle: MLA1 (no survival in exposed or control individuals).

### Bioassay 2


We found significant differences in survival between La Balandra and Pinamar populations exposed to Bd and their respective controls ( $p < 0.001$ ); MST were 3.1 d and 5.0 d respectively. In specimens from the Pinamar population, survival decreased from 100% on day 4 to

0% on day 5, while for La Balandra the decrement was more gradual (Fig. 1b). All tadpoles in control groups survived throughout the bioassay. The survival analysis showed significant differences between La Balandra and Pinamar ( $p = 0.001$ ), and neither group had survivors at the end of the bioassay (day 5 for Pinamar and day 7 for La Balandra). Oral discs were deformed in about half of the larvae, including the total loss of keratinised mouth parts in the upper and/or lower jaws (Fig. 3).

We identified Bd thalli in mouthparts of individuals from La Balandra (6/10 infected individuals) and Pinamar (3/10), although infection was mild in both (1–10 zoosporangia), and negative in controls. Infection was detected in tadpoles with normal mouthparts and those with some degree of depigmentation. Histological analyses were negative for individuals from Pinamar and controls, whereas Bd sporangia were present in individuals from La Balandra (Fig. 4).


### Bioassay 3

Survival differed significantly between *P. fernandae* tadpoles inoculated with isolate JEL423 (MST=6.5 d), isolate MLA1 (MST=3.3 d), and control groups (100% survival;  $p < 0.001$ ), as well as between the treatment groups (Fig. 1C;  $p = 0.003$ ). Between 50 and 90% of treated larvae and controls presented loss of keratinised mouth parts (typified in Bioassay 2). Bd thalli were detected in tadpoles exposed to either Bd strain in mouthparts either with or without deformation (6/10 and 2/10 infected individuals for MLA1 and JEL423, respectively), but not in control tadpoles. Histological analysis yielded negative Bd infection for both treatments and controls. The contingency table with all individuals of *P. fernandae* showed a dependency between depigmentation/no Bd (19/40) and normal/no Bd (2/40). Although no formal behavioural analyses were performed, we observed a change in the normal activity of the exposed tadpoles (Bioassays 1, 2 and 3) such as slow reaction to stimuli.



**Fig. 2.** *Rhinella fernandae* fresh oral disc surface (without stain), infected with Bd (400X). Note mature and empty zoosporangium in the stratum corneum (arrow). Scale bar=10 µm.





**Fig. 3.** Normal (A; 25 stage; Pinamar population) and abnormal (B; 38 stage; La Balandra population) oral discs in *Physalaemus fernandezae* tadpoles (Gosner, 1960). Arrows indicate lower sheath jaw (A) and lower jaw (B; showing lack of keratinised structures). Scale bars=200 µm (A); 500 µm (B).

## DISCUSSION


To our knowledge, our results represent the first report of disease susceptibility in native South American tadpoles experimentally exposed to *Bd*.

### Bioassay 1

All the species we tested died after exposure to *Bd* but varied widely in their sensitivity; whereas all individuals of *R. fernandezae* and *S. squalirostris* died on day 2, other species survived between 8 and 14 days. This low larval survival, recorded also in other species (Blaustein et al., 2005; Garner et al., 2009; Paetow et al., 2013; Hanlon & Parris, 2014), shows that amphibian larvae not only act as carriers and reservoirs of *Bd* (Narayan et al., 2014) but also experience mortality. The high mortality and low infection degree registered suggests that the study species are sensitive to exposure of *Bd*. That *Bd*-exposed tadpoles could invest a high amount of energy to prevent

infection, perhaps through mechanisms that inhibit zoospore attachment to host cells, could ultimately lead to larval mortality before metamorphosis (Garner et al., 2009). High concentration of *Bd* zoospores also produce harmful chemicals which might have contributed to the rapid mortality observed in our experiments (McMahon et al., 2012).

The absence of oral disc deformation in all species except *P. fernandezae* may be linked to a short time of exposure. As evidenced by the examination of unstained mouthparts, *Bd* was present in all species except *L. latrans* (see Peterson et al., 2007 for a similar example on another species). Although tadpoles of *L. latrans* may be resistant, the presence of *Bd* was confirmed in adults and juveniles of wild populations (Herrera et al., 2005; Ghirardi et al., 2009). Given the localised nature of an early stage of infection (Berger et al., 1999), false negatives could arise from histological analyses. This can be related to the absence of mouthpart deformation,



**Fig. 4.** Oral disc sections of *Physalaemus fernandezae* tadpoles from La Balandra population showing empty *Batrachochytrium dendrobatidis* zoosporangia (arrow). Scale bar=14 µm.



provided there exists a strong association between oral disc depigmentation and histological confirmation of *Bd* (Rachowicz & Vredenburg, 2004; Knapp & Morgan, 2006). The presence of deformations on the oral disc of *P. fernandezae* is noteworthy, and is discussed below.

### Bioassay 2

Although the survival time of tadpoles from both *P. fernandezae* populations was similar, individuals from Pinamar died in large numbers a single day, a difference which may have been caused by genetic differences between the two groups. Natale (2006) found differences in the sensitivity of tadpoles to a Chromium (VI) solution in two populations of *P. fernandezae*, but studies on mortality variation among different tadpole populations exposed to *Bd* are largely lacking (for examples on adults see Tobler & Schmidt, 2010; Bradley et al., 2015; Piovio-Scott et al., 2015). Although no individuals from either population survived longer than 7 days, survival in La Balandra tadpoles began to decrease 3 days earlier than for Pinamar. Individuals from Pinamar population were on average at earlier developmental stages compared to La Balandra when exposed to *Bd*, which might lead to a higher sensitivity to infection (*Bd*: Hanlon & Parris, 2014; see also Bunn et al., 2001; Johnson et al., 2011).

The pattern of mouthpart deformation in both the treatment and control groups was more consistent with that shown by tadpoles exposed to low temperatures (6°C) than with the pattern of 'gaps' that characterises *Bd* infection (Rachowicz & Vredenburg, 2004). We also identified *Bd* sporangia in both normal and abnormal mouthparts, suggesting loss of keratinised mouthparts in the absence of *Bd*, as well as unaffected mouthparts in the presence of *Bd* (see also Blaustein et al., 2005; Padgett-Flohr & Goble, 2007; Smith & Weldon, 2007). It is worth considering that *P. fernandezae* tadpoles generally have abnormalities in oral disc structures and in the pattern of ossification when reared under laboratory conditions (Barrasso, unpublished).

### Bioassay 3

Survival of groups inoculated with the Argentina (MLA1) and Panama (JEL423) *Bd* strains differed markedly. While the survival of both groups declined 48 hours after exposure, all individuals exposed to the Argentina strain died twice as fast than individuals exposed to the Panama strain. Experiments comparing the effects of different *Bd* strains revealed differences among host species which may be associated with environmental factors (Berger et al., 2005; Retallick & Miera, 2007; Gahl et al., 2012). JEL423 was isolated 6 years before MLA1 from an adult of *A. lemur* from Panama, and has produced symptoms and mortality in different species (Becker & Harris, 2010; Brannelly et al., 2012), whereas MLA1 was isolated from larvae of *H. cordobae* from a mountain stream in San Luis province (Argentina), and these are the first bioassays performed with this strain. Differences in the *in vitro* handling of the strains as well as in the time since their isolation may cause changes in their virulence (Berger et al., 2005; Brem et al., 2013). MLA1 has larger sporangia than JEL423 (Arellano et al., 2010), supporting its higher

virulence (see also Fisher et al., 2009). Genomic studies have revealed deep phylogenetic diversity, cryptic recombination and the existence of *Bd*-specific genes with possible pathogenicity factors (Joneson et al., 2011; Farrer et al., 2013; Rosenblum et al., 2013). The two strains used in our experiment are included in a global panzootic lineage (GPL) that contains the most infectious *Bd* isolates (Lips et al., 2006; Becker & Harris, 2010; Brannelly et al., 2012; Gahl et al., 2012; Rosenblum et al., 2013). Different effects on the survival of tadpoles can also be attributed to immunotoxicity (Piovio-Scott et al., 2015).

The finding that prevails in all experiments was high mortality of tadpoles, although reports of mass mortalities in the wild are lacking (but see Barrionuevo & Magione, 2006; Ghirardi et al., 2014). Tadpoles were exposed to concentrations of *Bd* zoospores ( $6 \times 10^4 \text{ ml}^{-1}$ ) which are likely higher than concentration in nature where *Bd* diffuses and becomes reduced through predators that forage on *Bd* zoospores (Searle et al., 2013; Schmeller et al., 2014; Groner & Relyea, 2015), and where chemical agents can have a fungicidal effect (Gahl et al., 2011; Hanlon & Parris, 2014; Rumschlag et al., 2014). Native tadpoles also might only experience *Bd*-caused declines when subjected to more virulent or allopatric strains (James et al., 2009).

## ACKNOWLEDGEMENTS

We thank S.M. Arellano for help editing the manuscript in English, J.E. Longcore for providing strain JEL423, and J. Grosso for providing the photo of *P. fernandezae* without infection. D.A.B. thanks L. Conte for field work assistance. D.A.B. and M.L.A. acknowledge Dirección de Flora y Fauna and Dirección de Contralor y Uso de los Recursos Naturales y Pesqueros de la Provincia de Buenos Aires, for granting Permit N° 190 and Expte. N° 22228-30/06 to conduct the study. This study was funded by Consejo Nacional de Investigaciones Científicas y Técnicas and Universidad Nacional de La Plata and partially supported by the Agencia Nacional de Promoción Científica y Tecnológica (PICT 2718, PICT 510, and PICT 2012-2315) and the Consejo Nacional de Investigaciones Científicas y Técnicas (PIP 11220120100510).

## REFERENCES

- Altig, R. (2007). Comments on the descriptions and evaluations of tadpole mouthpart anomalies. *Herpetological Conservation & Biology* 2, 1–4.
- Arellano, M.L., Marano, A.V., Natale, G.S., Steciow, M.M. & Lavilla, E.O. (2010). Aislamiento y caracterización morfológica de la primera cepa de *Batrachochytrium dendrobatidis* en Argentina. XI Congreso Argentino de Herpetología. Octubre de 2010. Buenos Aires, Argentina.
- Barrionuevo, S. & Mangione, S. (2006). Chytridiomycosis in two species of *Telmatobius* (Anura: Leptodactylidae) from Argentina. *Diseases of Aquatic Organisms* 73, 171–174.
- Becker, M.H. & Harris, R.N. (2010). Cutaneous bacteria of the redback salamander prevent morbidity associated with a lethal disease. *PLoS ONE* 5, e10957.

- Berger, L., Speare, R., Daszak, P., Green, D.E. et al. (1998). Chytridiomycosis causes amphibian mortality associated with population declines in the rain forests of Australia and Central America. *Proceedings of the National Academy of Sciences of the United States of America* 95, 9031–9036.
- Berger, L., Speare, R. & Kent, A. (1999). Diagnosis of chytridiomycosis in amphibians by histologic examination. Available from: <<http://www.jcu.edu.au/school/phtm/PHTM/frogs/histo/chhisto.htm>>. Accessed: March 1<sup>st</sup> 2014.
- Berger, L., Marantelli, G., Skerratt, L.F. & Speare, R. (2005). Virulence of the amphibian chytrid fungus *Batrachochytrium dendrobatidis* varies with the strain. *Diseases of Aquatic Organisms* 68, 47–50.
- Blaustein, A.R., Romansic, J.M., Scheessele, E.A., Han, B.A., et al. (2005). Interspecific variation in susceptibility of frog tadpoles to the pathogenic fungus *Batrachochytrium dendrobatidis*. *Conservation Biology* 19, 1–9.
- Bradley, P.W., Gervasi, S.S., Hua, J., Cothran, R.D., et al. (2015). Differences in sensitivity to the fungal pathogen *Batrachochytrium dendrobatidis* among amphibian populations. *Conservation Biology* 29, 1347–1356.
- Brannnelly, L.A., Richards-Zawacki, C.L. & Pessier, A.P. (2012). Clinical trials with itraconazole as a treatment for chytrid fungal infections in amphibians. *Diseases of Aquatic Organisms* 101, 95–104.
- Brem, F.M., Parris, M.J. & Padgett-Flohr, G.E. (2013). Re-Isolating *Batrachochytrium dendrobatidis* from an amphibian host increases pathogenicity in a subsequent exposure. *PLoS ONE* 8, e61260.
- Bunn, T.L., Parsons, P.J., Kao, E. & Dietert, R.R. (2001). Exposure to lead during critical windows of embryonic development: Differential immunotoxic outcome based on stage of exposure and gender. *Toxicological Sciences* 64, 57–66.
- Farrer, R.A., Henk, D.A., Garner, T.W.J., Balloux, F., et al. (2013). Chromosomal copy number variation, selection and uneven rates of recombination reveal cryptic genome diversity linked to pathogenicity. *PLoS Genetics* 9, e1003703.
- Fisher, M.C., Bosch, J., Yin, Z., Stead, D.A., et al. (2009). Proteomic and phenotypic profiling of the amphibian pathogen *Batrachochytrium dendrobatidis* shows that genotype is linked to virulence. *Molecular Ecology* 18, 415–429.
- Gahl, M.K., Pauli B.D. & Houlahan J.E. (2011). Effects of chytrid fungus and a glyphosate-based herbicide on survival and growth of wood frogs (*Lithobates sylvaticus*). *Ecological Applications* 21, 2521–2529.
- Gahl, M.K., Longcore, J.E. & Houlahan, J.E. (2012). Varying responses of Northeastern North American amphibians to the chytrid pathogen *Batrachochytrium dendrobatidis*. *Conservation Biology* 26, 135–141.
- Garner, T.W.J., Walker, S., Bosch, J., Leech, S., et al. (2009). Life history tradeoffs influence mortality associated with the amphibian pathogen *Batrachochytrium dendrobatidis*. *Oikos* 118, 783–791.
- Gervasi, S.S., Gondhalekar, C., Olson, D.H. & Blaustein, A.R. (2013). Host identity matters in the amphibian-*Batrachochytrium dendrobatidis* system: fine-scale patterns of variation in responses to a multi-host pathogen. *PLoS ONE* 8: e54490.
- Ghirardi, R., Lescano, J.N., Longo, M.S., Robledo, G., et al. (2009). *Batrachochytrium dendrobatidis* in Argentina: first record in *Leptodactylus gracilis* and another record in *Leptodactylus ocellatus*. *Herpetological Review* 40, 175–176.
- Ghirardi, R., Levy, M.G., López, J.A., Corbalán, V., et al. (2014). Endangered amphibians infected with the chytrid fungus *Batrachochytrium dendrobatidis* in Austral temperate wetlands from Argentina. *Herpetological Journal* 24, 129–133.
- Gosner, K.L. (1960). A simplified table for staging anuran embryos and larvae with notes on identification. *Herpetologica* 16, 183–190.
- Groner, M.L. & Relyea, R.A. (2015). Predators reduce *Batrachochytrium dendrobatidis* infection loads in their prey. *Freshwater Biology* 60, 1699–1704.
- Hanlon, S.M. & Parris, M.J. (2014). The interactive effects of chytrid fungus, pesticides, and exposure timing on gray treefrog (*Hyla versicolor*) larvae. *Environmental Toxicology Chemistry* 33, 216–222.
- Herrera, R., Steciow, M.M. & Natale, G.S. (2005). Chytrid fungus parasitizing the wild amphibian *Leptodactylus ocellatus* (Anura: Leptodactylidae) in Argentina. *Diseases of Aquatic Organisms* 64, 247–252.
- James, T.Y., Litvintseva, A.P., Vilgalys, R., Morgan, J.A.T. et al. (2009) Rapid global expansion of the fungal disease chytridiomycosis into declining and health amphibian populations. *PLoS Pathogens* 5, 1–12.
- Johnson, P.T.J., Kellermanns, E. & Bowerman, J. (2011). Critical windows of disease risk: Amphibian pathology driven by developmental changes in host resistance and tolerance. *Functional Ecology* 25, 726–734.
- Joneson, S., Stajich, J.E., Shiu, S-H. & Rosenblum, E.B. (2011). Genomic transition to pathogenicity in chytrid fungi. *PLoS Pathogens* 7, e1002338.
- Knapp, R.A. & Morgan, J.T. (2006). Tadpole mouthpart depigmentation as an accurate indicator of chytridiomycosis, an emerging disease of amphibians. *Copeia* 2, 188–197.
- Kielgast, J., Rödder, D., Veith, M. & Lötters, S. (2009) Widespread occurrence of the amphibian chytrid fungus in Kenya. *Animal Conservation* 13, 1–8.
- Langhammer, P.F., Burrowes, P.A., Lips, K.R., Bryant, A.B. & Collins, J.P. (2014). Susceptibility to the amphibian chytrid fungus varies with ontogeny in the direct-developing frog, *Eleutherodactylus coqui*. *Journal of Wildlife Disease* 50, 438–446.
- Lips, K.R., Brem, F., Brenes, R., Reeve, J.D., et al. (2006). Emerging infectious disease and the loss of biodiversity in a Neotropical amphibian community. *Proceedings of the National Academy of Sciences of the United States of America* 103, 3165–3170.
- Longcore, J.C., Pessier, A.P. & Nichols, D.K. (1999). *Batrachochytrium dendrobatidis* gen. Et sp. nov., a chytrid pathogenic to amphibians. *Mycologia* 91, 219–227.
- McMahon, T.A., Brannnelly, L.A., Chatfield, M.W.H., Johnson, P.T.J. & Joseph, M.B. et al. (2013). Chytrid fungus *Batrachochytrium dendrobatidis* has nonamphibian hosts and releases chemicals that cause pathology in the absence of infection. *Proceedings of the National Academy of Sciences of the United States of America* 110, 210–215.
- Mitchell, K., Churcher, T., Garner, T. & Fisher, M. (2008). Persistence of the emerging pathogen *Batrachochytrium dendrobatidis* outside the amphibian host greatly increases the probability of host extinction. *Proceedings of the Royal Society B: Biological Sciences* 275, 329–334.

- Narayan, E.J., Graham, C., McCallum, H. & Hero, J-M (2014). Over-wintering tadpoles of *Mixophyes fasciolatus* act as reservoir host for *Batrachochytrium dendrobatidis*. *PLoS ONE* 9, e92499.
- Natale, G.S. (2006). *Analysis of a community of ecotoxicological anurans of the Pampas. Effect of Cr (VI) on embryonic and larval different species of taxocomunidad* [in Spanish]. Doctoral Thesis. Facultad de Ciencias Naturales y Museo. Universidad Nacional de La Plata.
- Ortiz-Santaliestra, M.E., Rittenhouse, T.A.G., Cary, T.L. & Karasov, W.H. (2013). Interspecific and postmetamorphic variation in susceptibility of three North American anurans to *Batrachochytrium dendrobatidis*. *Journal of Herpetology* 47, 286–292.
- Padgett-Flohr, G.E. & Goble, M.E. (2007). Evaluation of tadpole mouthparts depigmentation as a diagnostic test for infection by *Batrachochytrium dendrobatidis* for four California anurans. *Journal of Wildlife Diseases* 43, 600–699.
- Paetow, L.J., McLaughlin, J.D., Pauli, B.D. & Marcogliese, D.J. (2013). Mortality of American bullfrog tadpoles *Lithobates catesbeianus* infected by *Gyrodactylus jennyae* and experimentally exposed to *Batrachochytrium dendrobatidis*. *Journal of Aquatic Animal Health* 25, 15–26.
- Peterson, J.D., Wood, M.B., Hopkins, W.A., Unrine, J.M. & Mendonça, M.T. (2007). Prevalence of *Batrachochytrium dendrobatidis* in American Bullfrog and Southern Leopard Frog larvae from wetlands on the Savannah River Site, South Carolina. *Journal of Wildlife Disease* 43, 450–460.
- Piovia-Scott, J., Pope K., Worth S.J., Rosenblum, E.B., et al. (2015). Correlates of virulence in a frog-killing fungal pathogen: evidence from a California amphibian decline. *The ISME Journal* 9, 1570–1578.
- Rachowicz, L.J. & Briggs, C.J. (2007) Quantifying the disease transmission function: effects of density on *Batrachochytrium dendrobatidis* transmission in the mountain yellow-legged frog *Rana muscosa*. *Journal of Animal Ecology* 76, 711–721.
- Rachowicz, L.J. & Vredenburg, V.T. (2004). Transmission of *Batrachochytrium dendrobatidis* within and between amphibian life stages. *Diseases of Aquatic Organisms* 61, 75–83.
- Reeder, N.M.M., Pessier, A.P. & Vredenburg, V.T. (2012). A reservoir species for the emerging amphibian pathogen *Batrachochytrium dendrobatidis* thrives in a landscape decimated by disease. Litvintseva AP, ed. *PLoS ONE* 7, e33567.
- Retallick, R.W.R. & Miera, V. (2007). Strain differences in the amphibian chytrid *Batrachochytrium dendrobatidis* and non-permanent, sub-lethal effects of infection. *Diseases of Aquatic Organisms* 75, 201–207.
- Rosenblum, E.B., James, T.Y., Zamudio, K.R., Poorten, T.J., et al. (2013). Complex history of the amphibian-killing chytrid fungus revealed with genome resequencing data. *Proceedings of the National Academy of Sciences of the United States of America* 110, 9385–9390.
- Rumschlag, S. L., Boone, M.D. & Fellers, G. (2014). The effects of the amphibian chytrid fungus, insecticide exposure, and temperature on larval anuran development and survival. *Environmental Toxicology and Chemistry* 33, 2545–2550.
- Ryan, M., Lips, K.R. & Eichholza, M.W. (2008). Decline and extirpation of an endangered Panamanian stream frog population (*Craugastor punctariolus*) due to an outbreak of chytridiomycosis. *Biological Conservation* 141, 1636–1647.
- Schmeller, D.S., Blooi, M., Martel, A., Garner T.W.J., et al. (2014). Microscopic aquatic predators strongly affect infection dynamics of a globally emerged pathogen. *Current Biology* 24, 176–180.
- Searle, C.L., Gervasi, S.S., Hua, J., Hammond, J.I., et al. (2011) Differential host susceptibility to *Batrachochytrium dendrobatidis*, an emerging amphibian pathogen. *Conservation Biology* 25, 965–974.
- Searle, C. L., Mendelson, J.R., Green L.E. & Duffy M.A. (2013). *Daphnia* predation on the amphibian chytrid fungus and its impacts on disease risk in tadpoles. *Ecology and Evolution* 3, 4129–4138.
- Smith, K.G. & Weldon, C. (2007). A conceptual framework for detecting oral chytridiomycosis in tadpoles. *Copeia* 4, 1024–1028.
- Spitzen-Van Der Sluijs, A., Martel, A., Hallmann, C.A., Bosman, W., et al. (2014). Environmental determinants of recent endemism of *Batrachochytrium dendrobatidis* infections in amphibian assemblages in the absence of disease outbreaks. *Conservation Biology* 28, 1302–1311.
- Symonds, E.P., Hines, H.B., Bird, P.S., Morton, J.M. & Mills, P.C. (2007). Surveillance for *Batrachochytrium dendrobatidis* using *Mixophyes* (Anura: Myobatrachidae) larvae. *Journal of Wildlife Diseases* 43, 48–60.
- Tobler, U. & Schmidt, B.R. (2010). Within- and among-population variation in chytridiomycosis-induced mortality in the toad *Alytes obstetricans*. *PLoS One* 5, e10927. doi:10.1371/journal.pone.0010927.
- Voyles, J., Rosenblum, E.B. & Berger, L. (2011). Interactions between *Batrachochytrium dendrobatidis* and its amphibian hosts: A review of pathogenesis and immunity. *Microbes and Infection* 13, 25–32.
- Vredenburg, V.T., Knapp, R.A., Tunstall, T.S. & Briggs, C.J. (2010). Dynamics of an emerging disease drive large-scale amphibian population extinctions. *Proceedings of the National Academy of Sciences of the United States of America* 107, 9689–9694.
- Woodhams, D.C., Alford, R.A. & Marantelli, G. (2003). Emerging disease of amphibians cured by elevated body temperature. *Diseases of Aquatic Organisms* 55, 65–67.

Accepted: 23 December 2015



